Grade Post Therapy Path (yp) (Lacrimal Sac)

Organization	Field Name	ID	Required
KCR	Grade Post Therapy Path (yp)	30138	yes
SEER	Grade Post Therapy Path (yp)	3845	yes

Note 1 Leave post therapy grade blank when

- No neoadjuvant therapy
- · Clinical or pathological case only
- Neoadjuvant therapy completed; surgical resection not done
- . There is only one grade available and it cannot be determined if it is clinical, pathological, post therapy clinical or post therapy pathological

Note 2 Assign the highest grade from the resected primary tumor assessed after the completion of neoadjuvant therapy.

Note 3 If there are multiple tumors with different grades abstracted as one primary, code the highest grade.

Note 4 Use the grade from the post therapy clinical work up from the primary tumor in different scenarios based on behavior or surgical resection

- Behavior
- Tumor behavior for the post therapy clinical and the post therapy pathological diagnoses are the same AND the post therapy clinical grade is the highest grade
- . Tumor behavior for post therapy clinical diagnosis is invasive, and the tumor behavior for the post therapy pathological diagnosis is in situ
- Surgical Resection
- Surgical resection is done of the primary tumor after neoadjuvant therapy is completed and there is no grade documented from the surgical resection
- · Surgical resection is done of the primary tumor after neoadjuvant therapy is completed and there is no residual cancer

Note 5 Code 9 (unknown) when

- · Surgical resection is done after neoadjuvant therapy and grade from the primary site is not documented
- Surgical resection is done after neoadjuvant therapy and there is no residual cancer
- · Grade checked "not applicable" on CAP Protocol (if available) and no other grade information is available

Code	Description
Α	Well differentiated
В	Moderately differentiated
С	Poorly differentiated
D	Undifferentiated, anaplastic
9	Grade cannot be assessed; Unknown
<blank></blank>	See Note 1