Grade Clinical (NET Colon and Rectum)

| Organization | Field Name | ID | Required |
|--------------|----------------|-------|----------|
| KCR | Grade Clinical | 30136 | yes |
| SEER | Grade Clinical | 3843 | yes |

Note 1 Grade Clinical must not be blank.

Note 2 Assign the highest grade from the primary tumor assessed during the clinical time frame.

Note 3 If there are multiple tumors with different grades abstracted as one primary, code the highest grade.

Note 4 Codes 1-3 take priority over codes A-D.

Note 5 Code 9 (unknown) when

- Grade from primary site is not documented
- Clinical workup is not done (for example, cancer is an incidental finding during surgery for another condition)
- Grade checked "not applicable" on CAP Protocol (if available) and no other grade information is available

Note 6 If there is only one grade available and it cannot be determined if it is clinical or pathological, assume it is a Grade Clinical and code appropriately per Grade Clinical categories for that site, and then code unknown (9) for Grade Pathological, and blank for Grade Post Therapy Clin (yc) and Grade Post Therapy Path (yp).

| Code | Description |
|------|---|
| 1 | G1: Mitotic count (per 10 HPF or 2mm2) less than 2 AND Ki-67 index (%) less than 3 Stated as WHO Grade 1 |
| 2 | G2: Mitotic count (per 10 HPF or 2mm2) equal 2-20 OR Ki-67 index (%) equal 3-20 Stated as WHO Grade 2 |
| 3 | G3: Mitotic count (per 10 HPF or 2mm2) greater than 20 OR Ki-67 index (%) greater than 20 Stated as WHO Grade 3 |
| Α | Well differentiated |
| В | Moderately differentiated |
| С | Poorly differentiated |
| D | Undifferentiated, anaplastic |
| 9 | Grade cannot be assessed (GX); Unknown |