Grade Pathological (Testis)

Organization	Field Name	ID	Required
KCR	Grade Pathological	30137	yes
SEER	Grade Pathological	3844	yes

Note 1 Grade Pathological must not be blank.

Note 2 Assign the highest grade from the primary tumor.

Note 3 If there are multiple tumors with different grades abstracted as one primary, code the highest grade.

Note 4 Use the grade from the clinical work up from the primary tumor in different scenarios based on behavior or surgical resection

Behavior

- · Tumor behavior for the clinical and the pathological diagnoses are the same AND the clinical grade is the highest grade
- · Tumor behavior for clinical diagnosis is invasive, and the tumor behavior for the pathological diagnosis is in situ

Surgical Resection

- · Surgical resection is done of the primary tumor and there is no grade documented from the surgical resection
- · Surgical resection is done of the primary tumor and there is no residual cancer

• No surgical resection

 Surgical resection of the primary tumor has not been done, but there is positive microscopic confirmation of distant metastases during the clinical time frame

Note 5 Code 9 (unknown) when

- Grade from primary site is not documented
- No resection of the primary site (see exception in Note 4, Surgical resection, last bullet)
- Neo-adjuvant therapy is followed by a resection (see Grade Post Therapy Path (yp))
- · Grade checked "not applicable" on CAP Protocol (if available) and no other grade information is available
- Clinical case only (see Grade Clinical)
- There is only one grade available and it cannot be determined if it is clinical, pathological, post therapy clinical or post therapy pathological

Code	Description	
Α	Well differentiated	
В	Moderately differentiated	
С	Poorly differentiated	
D	Undifferentiated, anaplastic	
9	Grade cannot be assessed; Unknown	